arXiv:1205.5151v1 [astro-ph.CO] 23 May 2012

Unimodular Constraint on global scale Invariance

Naveen K. Singh*

Theory Division, Physical Research Laboratory, Navrangpura, Ahmedabad - 380 009, India

Abstract: The global scale invariance along with the unimodular gravity in the vacuum is studied in this paper. The global scale invariant gravitational action which follows the unimodular general coordinate transformations is considered without invoking any scalar field. The possible solutions for the gravitational potential under linear field approximation for the allowed values of the introduced parameters of the theory are discussed. The modified solution has additional corrections along with the Schwarzschild solution. A comparative study of unimodular theory with conformal theory is also presented. Furthermore, the cosmological solution is studied and it is shown that the unimodular constraint preserve the de Sitter solution.

1 Introduction

Surprisingly, we live in the era of late time acceleration of universe [1, 2]. The acceleration in expansion of universe offers a new physics of unknown component " dark energy". The other matter, whose physical nature is partially known, "dark matter", is also one of the interesting subjects in cosmology. Dark matter was introduced by Fritz Zwicky in year 1933 [3, 4] to account for evidence of missing mass in Coma cluster. The more interesting issue to notice is that dark matter and dark energy dominate the energy density of the universe. The observed dark energy and dark matter contribute approximately 72.8% and 22.7% respectively to the total energy content of the universe. There are number of scalar field theories such as Quintessence, [5–10] K-essence [11–13], Chaplygin gas model [14–16] and theory of modified gravity such as $f(R)$ [17–22], DGP model [23, 24] etc. to describe the dark side of universe. In the recent literature [25–27], authors discuss the absence of the dark matter in the vicinity of solar system. The dark matter might be internal property of the space and hence might be explained by the modified theory of gravity [28–40]. In this paper, such a theory of modified gravity is considered which has global scale invariance along with the unimodular constraint to explain the galactic rotational curve. The scale invariance was first discussed by Weyl in 1929 [41]. This symmetry is broken if the theory contains any dimensionful parameters, such as particle masses, cosmological constant, gravitational constant etc.. One of the motivation for considering scale symmetry is that it might impose some constraints on the cosmological constant [42–47]. Hence it might help in solving its fine tuning problem. Although scale invariance is generally believed to be anomalous, it is possible to maintain this symmetry in the full quantum theory if the symmetry is broken by a soft mechanism [45–50]. This does not conflict with any known consequences of scale anomaly such as scale dependence of the coupling constants [45, 46]. It has been shown in many of the literatures [47, 49–53] that scale invariance has its advantage in explaining the dark sector of the universe. The local scale invariance explains the current era of the universe [51], favoring the ΛCDM model. In the papers [51–53] the symmetry breaking mechanism of local scale invariance generates the dark energy and vector field, which is gauge particle acts as dark matter field. In the literature [53], scale invariance has been discussed in details. The global scale transformation is given by

---

*email: naveenks@prl.res.in

$$\begin{array}{rcl} x^{\mu} & \rightarrow & \Lambda x^{\mu} \\ ds^2 & \rightarrow & \Lambda^2 ds^2 \end{array}$$

where $ds^2 = g_{\mu\nu}dx^{\mu}dx^{\nu}$ is the line element and Λ is a constant parameter. So we have

$$\bar{g}_{\mu\nu} \rightarrow \bar{g}_{\mu\nu} \tag{1}$$

The matter fields transform as follows,

$$\begin{array}{rcl} \phi & \rightarrow & \phi/\Lambda \\ A_{\mu} & \rightarrow & A_{\mu} \\ \psi & \rightarrow & \psi/\Lambda^{3/2} \end{array}$$

where ϕ, A_{μ} and ψ are scalar, vector and spinor fields. The above transformations make the action like $\int R^2\sqrt{-g}d^4x$ or $\int \phi^2 R\sqrt{-g}d^4x$ invariant.

The theories of higher order invariants in the action were initially started by Weyl in 1919 and by Eddington in 1923. The $f(R)$ theory has also been studied in the literatures [17–22,40,54–60]. One of the advantage of $f(R)$ theory is it describes the early universe [61,62]. Further, it also explains the late time acceleration [40,63] in the expansion of universe and might be an alternative for the dark matter [28–40]. The theory follows the principle of covariance similar as the Einstein-Hilbert action. It modifies the Einstein equation and so the solution for the gravitational potential. The only quadratic terms of the curvature scalar in the action preserves the global scale invariance. It is reasonable to take only quadratic terms of the curvature scalar in the action, since the solution could include the Schwarzschild solution and the corrections in addition [28]. Using the Gauss-Bonnet identity, we may write such action as following [64]

$$-\alpha_g \int (R^{\mu\nu}R_{\mu\nu} + \gamma R^2)\sqrt{-g}d^4x \tag{2}$$

It should be noted that the theory is now not conformal in general [28]. Conformal theory is one of the special case of (2), where $\gamma = -1/3$. In the section (2), the field equation of metric is derived. In the section 3, the corresponding equation for the unimodular theory is given. The field equation is solved and also the corresponding plot for galactic rotational curve is given in this section. The cosmological solution with unimodular constraint is discussed in the section (4). The last section (5) contains the discussions and the conclusions.

2 Field Equation

The variation of action 2 gives

$$\begin{aligned} \delta S &= -\alpha_g \int \Big[\sqrt{-g}(2\delta g^{\rho\mu}g^{\sigma\nu}R_{\rho\sigma}R_{\mu\nu} + 2g^{\rho\mu}g^{\sigma\nu}R_{\rho\sigma}\delta R_{\mu\nu} + 2\gamma R(\delta g^{\mu\nu}R_{\mu\nu} + g^{\mu\nu}\delta R_{\mu\nu})) \\ &- \frac{1}{2}\sqrt{-g}g_{\mu\nu}\left(\gamma R^2 + R_{\rho\sigma}R^{\rho\sigma}\right)\delta g^{\mu\nu}\Big]d^4x \end{aligned} \tag{3}$$

Now using the identity [65],

$$\delta R_{\mu\nu} = \left[(\delta\Gamma^{\lambda}_{\mu\nu})_{;\lambda} - (\delta\Gamma^{\lambda}_{\mu\lambda})_{;\nu} \right], \tag{4}$$

we may write the field equayion as

$$W^1_{\mu\nu} + W^2_{\mu\nu} = 0 \ . \tag{5}$$

Here $W^1_{\mu\nu}$ and $W^2_{\mu\nu}$ are the terms corresponding to the variation of R^2 and $R_{\rho\sigma}R^{\rho\sigma}$ and these are given as

$$\begin{aligned} W^1_{\mu\nu} &= -\frac{\gamma}{2}R^2 g_{\mu\nu} + 2\gamma\left[g^{\alpha\beta}R_{;\alpha;\beta}g_{\mu\nu} - R_{;\mu;\nu}\right] + 2\gamma R R_{\mu\nu}, \\ W^2_{\mu\nu} &= -\frac{1}{2}R_{\rho\sigma}R^{\rho\sigma}g_{\mu\nu} + 2R_{\mu\rho}R_{\nu\sigma}g^{\rho\sigma} - 2(R^{\alpha\beta})_{;\mu;\beta}g_{\alpha\nu} + (R_{\mu\nu})^{;\rho}_{;\rho} + (R^{\alpha\beta})_{;\beta;\alpha}g_{\mu\nu} \end{aligned} \tag{6}$$

respectively.

3 Unimodular Gravity

Unimodular gravity has been studied in the several literatures [66–80]. The theory is subclass of general theory of relativity but with a constraint in addition, i.e., the determinant of the metric is not dynamical; $g_{\mu\nu}\delta g^{\mu\nu} = 0$. The motivation of the unimodular gravity is to solve the cosmological constant problem as we don't have any such term in the action. Further, in the reference [66], authors discuss dynamics of expansion of universe with the unimodular theory of gravity taking the dynamical part of determinant of metric as a separate scalar field. However, in this paper, any scalar field is not considered. The condition $g_{\mu\nu}\delta g^{\mu\nu} = 0$ modifies the Einstein equation as following [67, 69],

$$R_{\mu\nu} - \frac{1}{4}g_{\mu\nu}R = \kappa\left(T_{\mu\nu} - \frac{1}{4}g_{\mu\nu}T\right) \ , \tag{7}$$

where κ is coupling constant, $T_{\mu\nu}$ is energy momentum tensor of source field and T is its trace. The Eq. 7 is traceless part of the Einstein equation. The variation of action 2 gives following field equation

$$\begin{aligned} W_{\alpha\beta} &= -\frac{1}{2}g_{\alpha\beta}\left(\gamma R^2 + R_{\rho\sigma}R^{\rho\sigma}\right) + 2\gamma\left[g^{\mu\nu}(R)_{;\mu;\nu}g_{\alpha\beta} - (R)_{;\alpha;\beta}\right] + 2R_{\alpha\nu}R_{\beta\rho}g^{\rho\nu} \\ &- 2\left(R^{\mu\nu}\right)_{;\alpha;\nu}g_{\mu\beta} + \left(R^{\mu\nu}\right)^{;\lambda}_{;\lambda}g_{\mu\alpha}g_{\nu\beta} + \left(R^{\mu\nu}\right)_{;\nu;\mu}g_{\alpha\beta} + 2\gamma R R_{\alpha\beta} = 0 \ . \end{aligned} \tag{8}$$

The same procedure of constraint of unimodular gravity over the action given in Eq. 2 gives the following field equation,

$$\begin{aligned} W^{uni}_{\alpha\beta} &= -\frac{1}{2}g_{\alpha\beta}\left(\gamma R^2 + R_{\rho\sigma}R^{\rho\sigma}\right) + 2\gamma\left[g^{\mu\nu}(R)_{;\mu;\nu}g_{\alpha\beta} - (R)_{;\alpha;\beta}\right] + 2R_{\alpha\nu}R_{\beta\rho}g^{\rho\nu} \\ &- 2\left(R^{\mu\nu}\right)_{;\alpha;\nu}g_{\mu\beta} + \left(R^{\mu\nu}\right)^{;\lambda}_{;\lambda}g_{\mu\alpha}g_{\nu\beta} + \left(R^{\mu\nu}\right)_{;\nu;\mu}g_{\alpha\beta} + 2\gamma R R_{\alpha\beta} - \frac{W}{4}g_{\alpha\beta} = 0 \ , \end{aligned} \tag{9}$$

where $W = W^\alpha_\alpha$ is trace of tensor $W_{\alpha\beta}$. The Eq. 9 is the traceless part of Eq. 8. Here $g_{\mu\nu}\delta g^{\mu\nu} = 0$ is used, i.e., the action does not have any constant term.

3.1 Vacuum Solution for the Conformal Theory

For $\gamma = -1/3$, W^{rr} component of Eq. (9) gives the following equation

$$\begin{aligned} B^{-1}W^{rr} &= \frac{1}{6}B'B''' - \frac{1}{12}B''^2 - \frac{1}{3r}\left(BB''' - B'B''\right) - \frac{1}{3r^2}\left(BB'' + B'^2\right) \\ &+ \frac{2}{3r^3}BB' - \frac{B^2}{3r^4} + \frac{1}{3r^4}\ , \end{aligned} \tag{10}$$

where the metric is given by,

$$ds^2 = -B(r)dt^2 + \frac{1}{B(r)}dr^2 + r^2 d\Omega\ . \tag{11}$$

The exact vacuum of Eq. (10) may be written as [28]

$$B(r) = 1 - \frac{C_1(2 - 3C_1C_2)}{r} - 3C_1C_2 + C_2 r - C_3 r^2\ , \tag{12}$$

where, C_1, C_2 and C_3 are constants.

3.2 Vacuum Solution for Unimodular Gravity

In this section, we solve for the gravitational potential with unimodular gravity considering the line element (11). $-(1/B)$ of $t-t$ component, $-B$ of $r-r$ component and $1/r^2$ of $\theta-\theta$ component of field Eq. 9 are given by

$$\begin{aligned} &(1+2\gamma)\frac{B''^2}{4} - \gamma\frac{B'^2}{r^2} + (1+4\gamma)\frac{B'B''}{r} - \frac{(1+2\gamma)}{r^4} - (1+4b)\frac{B^2}{r^4} \\ &+(2+6\gamma)\frac{B}{r^4} - (2+4\gamma)\frac{BB'}{r^3} + \gamma\frac{B'B'''}{2} - (2+3\gamma)\frac{BB'''}{r} \\ &-\gamma\frac{BB''}{r^2} - (1+\gamma)\frac{BB''''}{2} + 2(1+3\gamma)\frac{B'}{r^3} = 0\ , \end{aligned} \tag{13}$$

$$\begin{aligned} &-(1+2\gamma)\frac{B''^2}{4} + \gamma\frac{B'^2}{r^2} - (1+4\gamma)\frac{B'B''}{r} + \frac{(1+2\gamma)}{r^4} - (7+20\gamma)\frac{B^2}{r^4} \\ &+(6+18\gamma)\frac{B}{r^4} + (2+4\gamma)\frac{BB'}{r^3} - \gamma\frac{B'B'''}{2} - (2+5\gamma)\frac{BB'''}{r} \\ &+(4+13\gamma)\frac{BB''}{r^2} - (1+3\gamma)\frac{BB''''}{2} - 2(1+3\gamma)\frac{B'}{r^3} = 0 \end{aligned} \tag{14}$$

and

$$\begin{aligned} &-(1+2\gamma)\frac{B''^2}{4} + \gamma\frac{B'^2}{r^2} - (1+4\gamma)\frac{B'B''}{r} + \frac{(1+2\gamma)}{r^4} - (3+8\gamma)\frac{B^2}{r^4} \\ &+(2+6\gamma)\frac{B}{r^4} + (2+4\gamma)\frac{BB'}{r^3} - \gamma\frac{B'B'''}{2} - \gamma\frac{BB'''}{r} \\ &+(2+7\gamma)\frac{BB''}{r^2} - \gamma\frac{BB''''}{2} - (2+6\gamma)\frac{B'}{r^3} = 0 \end{aligned} \tag{15}$$

respectively. Now, considering linear approximation, i.e., $B(r) \approx 1+\phi(r)$, we have following three equations

$$-\frac{2\gamma\phi}{r^4}+\frac{2\gamma\phi'}{r^3}-\frac{\gamma\phi''}{r^2}-\frac{(2+3\gamma)\phi'''}{r}-\frac{(1+\gamma)\phi''''}{2}=0\ , \tag{16}$$

$$-\frac{(8+22\gamma)\phi}{r^4}-\frac{2\gamma\phi'}{r^3}+\frac{(4+13\gamma)\phi''}{r^2}-\frac{(2+5\gamma)\phi'''}{r}-\frac{(1+3\gamma)\phi''''}{2}=0 \tag{17}$$

and

$$-\frac{(4+10\gamma)\phi}{r^4}-\frac{2\gamma\phi'}{r^3}+\frac{(2+7\gamma)\phi''}{r^2}-\frac{\gamma\phi'''}{r}-\frac{\gamma\phi''''}{2}=0 \tag{18}$$

respectively. The solution of Eq. 16 is given by

$$\phi = C_1 r^{\frac{-2\gamma}{1+\gamma}}+\frac{C_2}{r}+C_3 r+C_4 r^2\ . \tag{19}$$

Plugging this solution, either in Eq. 17 or 18, we get the same constraint over the constants,

$$(1+3\gamma)\Big[-(6\gamma^3+\gamma^2-5\gamma-2)C_1+2(1+\gamma)^4 r^{1+\frac{2\gamma}{1+\gamma}}C_3\Big]=0\ . \tag{20}$$

Now, we have different solution for allowed values of γ and other constants. The constraint Eq. 20 gives one of the case where $\gamma = -1/3$. For this value we get

$$\phi = \frac{C_2}{r}+(C_1+C_3)r+C_4 r^2\ , \tag{21}$$

which is same solution as in Eq. (12) for the conformal theory. However, for $\gamma \neq -1/3$, we have $C_3 = 0$ and

$$6\gamma^3+\gamma^2-5\gamma-2=0\ , \tag{22}$$

which implies

$$\gamma=-\frac{2}{3},\quad \gamma=-\frac{1}{2}\ \text{ and } \gamma = 1\ . \tag{23}$$

For these values of γ the solutions are given by

$$\phi = C_1 r^4+\frac{C_2}{r}+C_4 r^2, \tag{24}$$

$$= \frac{C_2}{r}+(C_1+C_4)r^2, \tag{25}$$

$$= \frac{(C_1+C_2)}{r}+C_4 r^2 \tag{26}$$

respectively. The solution 25 or 26 with the data of galactic rotational curve [81,82] given in table. (1) for the Milky Way galaxy is plotted in Fig 1. The effective velocity of star may be written as

$$v^2=\frac{f}{2}\left(r\frac{\partial\phi}{\partial r}\right)\ , \tag{27}$$

Table 1: Table for galactic rotational curve.

Distance in Kpc	Velocity v (100 Km/sec)	Error (100 Km/sec)
2.97	2.03	.225
3.82	2.16	.180
4.67	2.26	.185
5.52	2.18	.090
6.37	2.27	.040
7.22	2.25	.045
7.50	2.15	.200
8.07	2.21	.040
12.5	2.26	.200
17.5	1.80	.200
22.5	1.64	.200
27.5	1.83	.200
32.5	1.43	.220
37.5	1.83	.390
42.5	2.03	.350
47.5	1.66	.300
55.0	1.80	.350

where $f = 9 \times 10^6$ to make velocity unit as $(100 Km/sec)$. For the best fit, the values of constants are

$$\begin{aligned} C_2 = -7.39 \times 10^{-6}\ Kpc \ \text{ and } \ (C_1 + C_4) = 1.67 \times 10^{-10}\ Kpc^{-2} \ \text{ for solution (25),} \\ (C_1 + C_2) = -7.39 \times 10^{-6}\ Kpc \ \text{ and } \ C_4 = 1.67 \times 10^{-10}\ Kpc^{-2} \ \text{ for solution (26).} \end{aligned} \tag{28}$$

The plot is shown by the dotted line. The further plot of the solution (24) with solid line is shown in Fig. 1. The values of constants for this case are as following

$$C_1 = -5.16 \times 10^{-14}\ Kpc^{-4},\ C_2 = -7.22 \times 10^{-6}\ Kpc \ \text{ and } \ C_4 = 3.78 \times 10^{-10}\ Kpc^{-2}\ . \tag{29}$$

The plot for the conformal theory is also shown with dashed line, where the gravitational potential is given by the Eq. (12) and for the best fit the values of the constants are given by

$$C_1 = 2.6526 \times 10^{-6}\ Kpc,\ \ C_2 = 5.0460 \times 10^{-8}\ Kpc^{-1} \text{ and } \ C_3 = 4.1366 \times 10^{-10}\ Kpc^{-2}\ . \tag{30}$$

The values of χ^2_{min}[1] per degree of freedom for the best fit for the solution (12), (24) and (25) are given by 3.19, 5.54 and 6.15 respectively. However, the solution (24) gives the best fit for the scale $> 15 Kpc$ as shown in the Fig. (1).

---

[1]
$$\chi^2 = \Sigma_i \left(\frac{theo_i - expt_i}{error} \right)^2 \tag{31}$$



Figure 1: The variation of velocity with distance r. The dotted curve is plot of solution given by (25) or (26) whereas solid curve is for the solution (24). The dashed curve is for the solution (12). Data for Milky Way Galaxy is shown with the error bar.

4 Cosmological Solution

It is known to us that Gauss-Bonnet action explains acceleration in the expansion of the universe [83–85] . Further, in the modified theory of gravity $f(R) = R^2$, we have exact de-Sitter solution [40] for the vacuum. In this section, we test it explicitly as now the action has the unimodular constraint in addition. For the FRW metric $[-1, a^2, a^2, a^2]$, where a is scale factor of the universe, Eq. (9) gives the same equation for $0-0$ and $i-j$ components and it is given by

$$-(6+18\gamma)\left(\frac{a'}{a}\right)^4 + (9+27\gamma)\frac{a'^2a''}{a^3} - (3+9\gamma)\left(\frac{a''}{a}\right)^2 + (1+3\gamma)\frac{a'a'''}{a^2} - (1+3\gamma)\frac{a''''}{a} = 0 \; . \tag{32}$$

The Eq. (32) may be written as independent of the parameter γ as

$$-6\left(\frac{a'}{a}\right)^4 + 9\frac{a'^2a''}{a^3} - 3\left(\frac{a''}{a}\right)^2 + \frac{a'a'''}{a^2} - \frac{a''''}{a} = 0 \; . \tag{33}$$

Looking over Eq. (33), one may conclude for the exact de-Sitter solution,

$$a = a_0 e^{H_0 t} \; , \tag{34}$$

which explain the acceleration in the universe-expansion, where H_0 is Hubble constant.

5 Discussion and Conclusions

A scale invariant model of higher order invariant in the action is presented. The unimodular constraint on the theory is also considered. Scale invariance allows only quadratic terms of curvature scalar in the action, whereas consideration of unimodular theory in addition constrain on the values of the parameter of the resulting theory. It is shown that for the parameter $\gamma = -1/2$ and 1, the solution of the gravitational potential includes the Schwarzschild solution as well as the term corresponding to the integration constant. The solution for this case explains the galactic rotational curve, but the corresponding gravitational field increases as distance increases whereas for $\gamma = -2/3$, the solution has one more term proportional to r^4 so that the velocity or corresponding gravitational field decreases after $\sim 42\ Kpc$. Furthermore, the solution of conformal theory is recovered for $\gamma = -1/3$. The conformal solution has a lighter bump at $\sim 30\ Kpc$. Hence, the unimodular theory of gravity has good behavior for the large scale rather than that of conformal theory. The proper scale invariant matter source term in the action might describe the rotational curve for the low range. We will proceed it further in the future publication. The theory is interesting as it does not require the dark matter which has not been observed in the solar neighborhood so far. Furthermore, the de Sitter solution is also obtained for the considered theory explaining the dynamics of current era.

Acknowledgements
Author thanks Prof. Subhendra Mohanty, Prof. Pankaj Jain and Girish Chakrabarty for useful discussions.

References

[1] S. Perlmutter *et al.*, Astrophys. J. **517**, 565 (1999).

[2] A. G. Riess *et al*, Astron. J. **116**, 1009 (1998).

[3] M. S. Turner, Phys. Scr. Vol. **T36**, 167-182 (1991).

[4] J. Einasto, Astronomy and Astrophysics 2010, [Eds. Oddbjorn Engvold, Rolf Stabell, Bozena Czerny, John Lattanzio], in Encyclopedia of Life Support Systems (EOLSS), Developed under the Auspices of the UNESCO, Eolss Publishers, Oxford ,UK.

[5] C. Wetterich, Nucl. Phys. B **302**, 668 (1988).

[6] B. Ratra and P. J. E. Peebles, Phys. Rev. D **37**, 3406 (1988).

[7] Y. Fujii and T. Nishioka, Phys. Rev. D **42**, 361 (1990).

[8] T. Chiba, N. Sugiyama and T. Nakamura, MNRAS **289**, L5 (1997).

[9] S. M. Carroll, Phys. Rev. Lett. **81**, 3067 (1998).

[10] E. J. Copeland, M. Sami, S. Tsujikawa, Int. J. Mod. Phys. **D15**, 1753-1936 (2006).

[11] T. Chiba, T. Okabe and M. Yamaguchi, Phys. Rev. D **62**, 023511 (2000).

[12] C. Armendariz-Picon, V. Mukhanov, and P. J. Steinhardt, Phys. Rev. Lett. **85**, 4438 (2000).

[13] C. Armendariz-Picon, V. Mukhanov, and P. J. Steinhardt, Phys. Rev. D **63**, 103510 (2001).

[14] G. F. R. Ellis and W. Stoeger , Class. Quant. Grav. 4, 1697 (1987).

[15] A. Y. Kamenshchik, U. Moschella and V. Pasquier, Phys. Lett. B **511**, 265 (2001).

[16] N. Bilic, G. B. Tupper and R. D. Viollier, Phys. Lett. B **535**, 17 (2002).

[17] C. H. Brans and R. H. Dicke, Phys. Rev. **124**, 925 (1961);
P. Jordan, Z. Phys. **157**, 112-121 (1959).

[18] R. Utiyama and B. S. DeWitt, J. Math. Phys. **3**, 608 (1962).

[19] D. Lovelock, J. Math. Phys.**12**, 498 (1971).

[20] H. A. Buchdahl, Mon. Not. R. Astron. Soc. **150**, 1 (1970).

[21] M. Milgrom, Astrophys. J. **270**, 365 (1983).

[22] J. Bekenstein, Phys. Rev. D **70**, 083509 (2004).

[23] G. Dvali, G. Gabadadze and M. Porrati, Phys. Lett. B **485**, 208 (2000).

[24] A. Nicolis, R. Rattazzi and E. Trincherini, Phys. Rev. D **79**, 064036 (2009).

[25] C. Moni Bidin, G. Carraro, R. A. Mendez, R. Smith, Accepted for publication in the Astrophysical Journal, [arXiv:1204.3924v1].

[26] S. Garbari, J.I. Read, G. Lake, MNRAS **416**, 2318 (2011), [arXiv:1105.6339.].

[27] M. Creze, E. Chereul, O. Bienayme and C. Pichon, A & A, **329**, 920 (1998).

[28] P. D. Mannheim and D. Kazanas, Astrophysical Journal, **342**, p. 635-638 (1989).

[29] S. Capozziello, V. F. Cardone, S. Carloni, and A. Troisi, Phys. Lett. A **326**, 292 (2004).

[30] S. Capozziello, V. F. Cardone, S. Carloni, and A. Troisi, AIP Conf. Proc. **751**, 54 (2005).

[31] S. Capozziello, V. F. Cardone, and A. Troisi, Phys. Rev. D **71**, 043503 (2005).

[32] S. Capozziello, V. F. Cardone, and A. Troisi, JCAP **0608**, 001 (2006).

[33] S. Capozziello, V. F. Cardone, and A. Troisi, Mon. Not. Roy. Astron. Soc. **375**, 1423 (2007).

[34] L. Iorio and M. L. Ruggiero, Int. J. Mod. Phys. A **22**, 5379 (2007).

[35] S. Jhingan, et al., Phys. Lett. B **663**, 424 (2008), [eprint 0803.2613].

[36] C. F. Martins and P. Salucci, Mon. Not. Roy. Astron. Soc. **381**, 1103 (2007).

[37] S. Nojiri and S. D. Odintsov, [eprint 0801.4843] (2008).

[38] R. Saffari and Y. Sobouti, Astron. Astroph. **472**, 833 (2007).

[39] H. Zhao and B. Li, 2008, eprint 0804.1588.

[40] T. P. Sotiriou and V. Faraoni, Rev.Mod.Phys. **82**, 451-497, (2010).

[41] H. Weyl, Z. Phys. **56**, 330 (1929). [Surveys High Energ. Phys. **5**, 261 (1986)].

[42] P. D. Mannheim, Gen.Rel.Grav. **22**, 289 (1990).

[43] P. D. Mannheim, College Park Astrophys., 0541-544 (1990).

[44] S. Deser, Annals Phys. **59**, 248-253 (1970).

[45] M. Shaposhnikov and D. Zenhausern, Phys. Lett. B **671**, 162 (2009). [arXiv:0809.3406 [hep-th]].

[46] M. Shaposhnikov and D. Zenhausern, Phys. Lett. B **671**, 187 (2009). [arXiv:0809.3395 [hep-th]].

[47] P. Jain, S. Mitra and N. K. Singh, JCAP **0803**, 011 (2008) [arXiv:0801.2041 [astro-ph]].

[48] F. Englert, C. Truffin and R. Gastmans, Nucl. Phys. B **117**, 407 (1976).

[49] P. Jain and S. Mitra, Mod. Phys. Lett. A **24**, 2069 (2009). [arXiv:0902.2525 [hep-ph]].

[50] P. K. Aluri, P. Jain, S. Mitra, S. Panda and N. K. Singh, Mod. Phys. Lett. A **25**, 1349 (2010). [arXiv:0909.1070 [hep-ph]].

[51] P. K. Aluri, P. Jain and N. K. Singh, Mod. Phys. Lett. A **24**, 1583 (2009). [arXiv:0810.4421 [hep-ph]].

[52] H. Cheng, Phys. Rev. Lett. **61**, 2182 (1988).

[53] H. Cheng and W. F. Kao, MIT preprint Print-88-0907 (1988).

[54] S. Capozziello and V. Faraoni, Fundamental Theories of Physics, Vol. **170**, Springer, (2010).

[55] S. Nojiri and S. D. Odintsov, Int. J. Geom. Meth. Mod. Phys. **4**, 115 (2007).

[56] S. Nojiri and S. D. Odintsov, Phys. Rep. **505**, 59 (2011).

[57] A. De Felice and S. Tsujikawa, Living Reviews in Relativity **13**: 3, (2010).

[58] S. Tsujikawa, Lect.Notes Phys.**800**: 99-145 (2010).

[59] S. Capozziello and M. De Laurentis, Phys. Rep. 509, 167-321 (2011).

[60] R. Ferraro, [arXiv:1204.6273 [gr-qc]].

[61] A. A. Starobinsky, Phys. Lett. B **91**, 99 (1980); Sov. Astron. Lett. **9**, 302 (1983).

[62] J. C. Hwang and H. Noh, Phys. Lett. B **506**, 13 (2001).

[63] S. Capozziello, S. Carloni, and A. Troisi, Recent Res. Dev. Astron. Astrophys. 1, 625 (2003).

[64] J. F. Donoghue, UMHEP-424, [arXiv:gr-qc/9512024v1].

[65] L. D. Landau and E. M. Lifshitz, The Classical Theory of Fields (Butterworth-Heinemann, 1980).

[66] P. Jain, P. Karmakar, S. Mitra, S. Panda and N. K. Singh, JCAP **05** 020 (2012).

[67] J.L. Anderson and D. Finkelstein, Am. J. Phys. **39**, 901 (1971).

[68] A. Einstein, in *The Principle of Relativity*, edited by A. Sommerfeld (Dover, New York, 1952).

[69] S. Weinberg, Rev. Mod. Phys. **61**, 1 (1989).

[70] J. J. van der Bij, H. van Dam and Y. J. Ng, Physica **116A**, 307 (1982).

[71] A. Zee, in *High Energy Physics: Proceedings of the 20th Annual Orbis Scientiae, Coral Gables, (1983)*, edited by B. Kursunoglu, S. C. Mintz, and A. Perlmutter (Plenum, New York, 1985).

[72] W. Buchmuller and N. Dragon, Phys. Lett. B **207**, 292 (1988).

[73] M. Henneaux and C. Teitelboim, Phys. Lett. B **222**, 195 (1989).

[74] W. G. Unruh, Phys. Rev. D **40**, 1048 (1989).

[75] Y. J. Ng and H. van Dam, J. Math. Phys. **32**, 1337 (1991).

[76] D. R. Finkelstein, A. A. Galiautdinov and J. E. Baugh, J. Math. Phys. **42**, 340 (2001) [arXiv:gr-qc/0009099].

[77] E. Alvarez, JHEP **0503**, 002 (2005) [arXiv:hep-th/0501146].

[78] E. Alvarez, D. Blas, J. Garriga, E. Verdaguer Nucl. Phys. B **756**, 148 (2006) [arXiv:hep-th/0606019].

[79] A. H. Abbassi and A. M. Abbassi, Class. Quant. Grav. **25**, 175018 (2008) [arXiv:0706.0451 [gr-qc]].

[80] G. F. R. Ellis, H. van Elst, J. Murugan and J. P. Uzan, [arXiv:1008.1196 (gr-qc)].

[81] X.X. Xue et al., Astrophys. J. **684**, 1143 (2008).

[82] M. Honma and Y. Sofue, PASJ **49**, 453 (1997).

[83] S. Nojiri, S. D. Odintsov, M. Sasaki, Phys. Rev. D **71**, 123509 (2005).

[84] S. Y. Zhou, E. J. Copeland, P. M. Saffin, JCAP **0907** 009 (2009).

[85] Y. Ito, S. Nojiri, Phys.Rev. D **79**, 103008 (2009).